

The Central African Journal of Medicine



Editor:

MICHAEL GELFAND, C.B.E., M.D., F.R.C.P.

Assistant Editor:

JOSEPH RITCHKEN, M.D.

Volume Nineteen

JANUARY - DECEMBER

1973

An investigation into the incidence and nature of Anaemia in pregnant African women

BY

J. D. KNOTTENBELT.

District Medical Officer, Bindura.

Report on the 6th year student project at the University of Rhodesia in the Department of Obstetrics and Gynaecology carried out in March, 1968.

INTRODUCTION

Anaemia is known to be prevalent among African women of the childbearing age groups in South and Central Africa, and it has been suggested by Davidson (1964) that iron deficiency is the commonest cause. The most likely causes of iron deficiency are dietary deficiency, and excessive loss due to childbearing and menstruation. Howard (1966), found that only 2.1 per cent. of 98 African schoolgirls aged 12½-17½ years had haemoglobin values less than 11 grams per 100 ml., with a mean value of 12.9. He found that parasitic infestations did not appear to be a major contributory cause of anaemia in his series. He remarked that these figures were in close agreement with those found by Beet (1949) in European school children. Donald (1964) states that in one series of women attending a Glasgow antenatal clinic, less than 50 per cent. of patients had a haemoglobin value greater than 10 G per cent., at their first visit. Accordingly an investigation into the incidence and nature of anaemias in African women attending Harare Hospital seemed relevant, especially as most patients come from the lower socio-economic groups.

METHODS

175 pregnant women attending professorial antenatal clinics for the first time during their current pregnancy were investigated. Their ages ranged from 16-45 years, their parities from 0-11, and the period of gestation from 16-38 weeks. They were questioned about all previous pregnancies, menstrual history and past medical and surgical history with special reference to any excess blood loss, blood transfusions, operations and abnormal pregnancy states such as abortion and ectopic pregnancy.

Using plastic disposable syringes, 12 ml of blood was withdrawn from each patient, 2 ml of which was placed in a plastic tube containing se-

questrene as anticoagulant, for haemoglobin, haematocrit and blood film examinations, and 10 ml of which was placed in a plastic test tube containing no anticoagulant. Plastic syringes and tubes were used since it has been shown that they are virtually iron-free.

The haemoglobin was estimated by a photometric cyanmethaemoglobin technique, and the packed cell volume (PCV) determined using a microhaematocrit centrifuge. By dividing the haemoglobin value in grams per cent. by the PCV, the mean corpuscular haemoglobin concentration per cent. was calculated (MCHC).

A thin blood film was made and stained with Leishman's stain. The red and white blood cell series were examined for any abnormality, with special reference to the changes of iron deficiency, hypochromic anaemia and macrocytic anaemia.

The serum iron was determined using a tripyridyl photometric technique. The serum iron of 10 male medical students was determined and the values obtained ranged from 90 to 150 µg per 100 ml., that is, within normal limits (50-150 µg.).

Haemoglobin value and serum iron value histograms were constructed. All patients were then divided into parity groups, and mean values for the haemoglobin, MCHC, and serum iron were calculated. The patients were then classified into 6 groups according to their haemoglobin, MCHC, and serum iron values, and blood film appearances.

Group I Patients with haemoglobin values over 10 G. per 100 ml., MCHC over 28 G.%, serum iron values over 50 µg per 100 ml., and a normal blood film.

Group II Patients with haemoglobin values over 10 G., normal MCHC., and blood film, but with serum iron less than 50 µg/100 ml.

Group III Patients with haemoglobin under 10 G./100 ml., MCHC less than 28 G.%, evidence of hypochromic anaemia on the blood film, and a low serum iron (less than 50 µg%).

Group IV Patients with haemoglobin under 10 g%, MCHC less than 28 G.%, evidence of hypochromic anaemia on the film, but with a serum iron above 50 µg/100 ml.

Group V Patients with haemoglobin under 10 g., normal MCHC, evidence of both hypochromic and macrocytic anaemia on the film and a low or low-normal serum iron, (all cases below 60 µg%).

Group VI Patients with a low haemoglobin, normal MCHC, evidence of macrocytic anaemia on the film, and a normal to high serum iron, (all cases had a value above 80 µg%).

RESULTS

(a) Haemoglobin distribution:

	Frequency						Percent of total.	
	0	10	20	30	40	50	60	
Less than 10 G %			20					11,4
10-10,9 G %				27				16,0
11-11,9 G %					49			28,1
12-12,9 G %						56		31,4
13-13,9 G %			20					11,4
14-14,9 G %	3							1,7

Mean haemoglobin = 11,6 G %
11,4 % have values below
10 G %

(b) Serum iron distribution:

	Frequency						Percent of total
Serum Iron in $\mu\text{g} \%$	0	10	20	30	40		
Less than 30		8					4,8
30-49			27				16,1
50-69				33			19,6
70-89					40		23,8
90-109				27			16,1
110-129			25				14,8
130-149	4						2,4
150+	4						2,4

Mean serum iron = 81 $\mu\text{g} \%$
20,9% have values below 50 $\mu\text{g} \%$

(c) Average values by parity:—

Parity	Haemoglobin	MCHC	Serum Iron ($\mu\text{g}\%$)	No. in sample
0	11,7	30,0	84	44
1	11,7	30,0	80	27
2	11,9	30,7	71	20
3	11,8	30,4	77	17
4	11,5	30,1	86	16
5	11,6	30,0	73	16
6	11,3	30,0	76	19
7	11,3	30,5	86	11
8	10,3	30,3	70	3
9	12,2	32,1	53	1
11	10,3	27,8	112	1
Total	11,6	30,2	81	175

(d) Classification by groups:

	No. in group	per cent. of total
Group I (Normal values)	129	73,2
Group II (Normal but iron low)	26	15,5
Group III (Hypochromic, iron low)	5	3,0
Group IV (Hypochromic, iron normal)	6	3,3
Group V (Mixed anaemia, iron low)	6	3,3
Group VI (Macrocytic, iron normal)	3	1,7

DISCUSSION

The haemoglobin histogram shows that 20 patients (11,4 per cent.) had a haemoglobin value less than 10 grams per 100 ml. This compares favourably with Scott's findings in Glasgow, and is no worse than values found in other studies of pregnant women in Britain. The value of 10 G% was chosen to make an adequate allowance for

physiological haemodilution in pregnancy, and this value seems to be fairly widely accepted in obstetrics. Of the 20 anaemic patients 11 had hypochromic anaemia, 6 had mixed hypochromic and macrocytic, and 3 had pure macrocytic anaemia. Thus 17 (9,7 per cent.) had evidence of hypochromic anaemia and 9 (5,1 per cent.) some evidence of possible folic acid deficiency.

The serum iron histogram shows that 35 (20,9 per cent.) patients, or approximately one in every five, had a serum iron below 50 $\mu\text{g}\%$. The limits of normal in European patients are taken as 50 and 150 $\mu\text{g}\%$ (100 $\mu\text{g}\%$ \pm 2 standard deviations). It is interesting to note that only 9 out of 35 patients with low serum iron values had significantly low haemoglobin values, although the average was lower than for the whole sample (10,9 G% — not statistically significant). 5 out of 8 patients with a serum iron of less than 30 $\mu\text{g}\%$, however, had a haemoglobin less than 10 G, with an average of 9,3 G (statistically significantly lower than the mean for all patients, 11,6 G%). Where serum iron values fell below 20 $\mu\text{g}\%$ all patients had a low haemoglobin value as well (average 7,8 G%). Thus while there is a general correlation between low serum iron and low haemoglobin values, it does not automatically follow that a patient with a low serum iron will be anaemic, or that a patient who is not anaemic has an adequate serum iron. This finding is consistent with the view that haemoglobin levels are maintained in preference to other iron stores in iron deficiency.

In the histories taken there was a remarkable lack of positive findings such as menorrhagia, abortions or ectopic pregnancy among those women found to be anaemic. Among those who had had such complications, the incidence of anaemia was no greater than that for the group as a whole. The abortion rate for all pregnancies in the women of the samples was 3,3 per cent., surprisingly lower than the expected 10-20 per cent. This may reflect a decreased tendency for those who abort to become pregnant again, or merely an increased interest in maintaining pregnancies in those women who attended this antenatal clinic.

In view of the high incidence of iron deficiency found in pregnant African women in this series, prophylactic iron therapy for every woman attending antenatal clinics would seem justified, regardless of the haemoglobin level found. Only four out of the series had a serum iron greater than 150 $\mu\text{g}\%$, so that the chances of iron overload do not seem to be very high. Since about 1 in 20 patients had either a mixed deficiency or a macrocytic anaemia, prophylactic folic acid

would appear to be indicated in those who are anaemic, especially as pernicious anaemia is very rare in African women of child bearing age. Serum folate determinations in a similar series would be most helpful in evaluating the true incidence of folic acid deficiency.

Contrary to what might have been expected, mean values for the haemoglobin, MCHC, and serum iron did not differ significantly on statistical analysis of the parity groups. Perhaps this is due to an average interval between pregnancies of 18 months to two years in the African multiparae in this series, or to a high dietary iron intake.

SUMMARY

(1) An account of an investigation into haemoglobin, MCHC and serum iron values in 175 African women attending Harare Hospital Antenatal Clinic for the first time in the current pregnancy is given.

(2) The mean haemoglobin for the group was 11,6 G%. 11,4% of the patients had values less

10 G%. 6,3 per cent. had evidence of hypochromic anaemia, 1,7 per cent macrocytic anaemia, and 3,3 per cent. had a mixed anaemia. The mean MCHC was 30,2 per cent.

(3) The mean serum iron was 81 μg per 100 ml. 20,9 per cent had levels below 50 μg %.

(4) There was no significant difference in haemoglobin, MCHC and serum iron values between groups of women of differing parity.

ACKNOWLEDGMENTS

I should like to thank Professor Philpott, Professor of the Department of Obstetrics and Gynaecology, University of Rhodesia, and Mr. A. Bedford, for their help to me in the planning and execution of the project.

REFERENCES

- DAVIDSON, L. A. G. (1964). *C. Afr. J. Med.* 10, 96. March, 1964.
HOWARD, J. K. (1966). *C. Afr. J. Med.* 12, 231. December, 1966.
DONALD, I. (1964). *Practical Obstetric problems.* Lloyd-Luke.

The Central African Journal of Medicine



Dr. DAVID LIVINGSTONE

Editor:

MICHAEL GELFAND, C.B.E., M.D., F.R.C.P.

Assistant Editor:

JOSEPH RITCHKEN, M.D.

Volume Nineteen

JANUARY - DECEMBER

1973

SALISBURY

RHODESIA

98 CEN IV

An investigation into the incidence and nature of Anaemia in pregnant African women

BY

J. D. KNOTTENBELT.

District Medical Officer, Bindura.

Report on the 6th year student project at the University of Rhodesia in the Department of Obstetrics and Gynaecology carried out in March, 1968.

INTRODUCTION

Anaemia is known to be prevalent among African women of the childbearing age groups in South and Central Africa, and it has been suggested by Davidson (1964) that iron deficiency is the commonest cause. The most likely causes of iron deficiency are dietary deficiency, and excessive loss due to childbearing and menstruation. Howard (1966), found that only 2.1 per cent. of 98 African schoolgirls aged 12½-17½ years had haemoglobin values less than 11 grams per 100 ml., with a mean value of 12.9. He found that parasitic infestations did not appear to be a major contributory cause of anaemia in his series. He remarked that these figures were in close agreement with those found by Beet (1949) in European school children. Donald (1964) states that in one series of women attending a Glasgow antenatal clinic, less than 50 per cent. of patients had a haemoglobin value greater than 10G per cent., at their first visit. Accordingly an investigation into the incidence and nature of anaemias in African women attending Harare Hospital seemed relevant, especially as most patients come from the lower socio-economic groups.

METHODS

175 pregnant women attending professorial antenatal clinics for the first time during their current pregnancy were investigated. Their ages ranged from 16-45 years, their parities from 0-11, and the period of gestation from 16-38 weeks. They were questioned about all previous pregnancies, menstrual history and past medical and surgical history with special reference to any excess blood loss, blood transfusions, operations and abnormal pregnancy states such as abortion and ectopic pregnancy.

Using plastic disposable syringes, 12 ml of blood was withdrawn from each patient, 2 ml of which was placed in a plastic tube containing se-

questrene as anticoagulant, for haemoglobin, haematocrit and blood film examinations, and 10 ml of which was placed in a plastic test tube containing no anticoagulant. Plastic syringes and tubes were used since it has been shown that they are virtually iron-free.

The haemoglobin was estimated by a photometric cyanmethaemoglobin technique, and the packed cell volume (PCV) determined using a microhaematocrit centrifuge. By dividing the haemoglobin value in grams per cent. by the PCV, the mean corpuscular haemoglobin concentration per cent. was calculated (MCHC).

A thin blood film was made and stained with Leishman's stain. The red and white blood cell series were examined for any abnormality, with special reference to the changes of iron deficiency, hypochromic anaemia and macrocytic anaemia.

The serum iron was determined using a tripyridyl photometric technique. The serum iron of 10 male medical students was determined and the values obtained ranged from 90 to 150 µg per 100 ml., that is, within normal limits (50-150 µg.)

Haemoglobin value and serum iron value histograms were constructed. All patients were then divided into parity groups, and mean values for the haemoglobin, MCHC, and serum iron were calculated. The patients were then classified into 6 groups according to their haemoglobin, MCHC, and serum iron values, and blood film appearances.

Group I Patients with haemoglobin values over 10 G. per 100 ml., MCHC over 28 G.%, serum iron values over 50 µg per 100 ml., and a normal blood film.

Group II Patients with haemoglobin values over 10 G., normal MCHC., and blood film, but with serum iron less than 50 µg/100 ml.

Group III Patients with haemoglobin under 10 G./100 ml., MCHC less than 28 G.%, evidence of hypochromic anaemia on the blood film, and a low serum iron (less than 50 µg%).

Group IV Patients with haemoglobin under 10 g.%, MCHC less than 28 G.%, evidence of hypochromic anaemia on the film, but with a serum iron above 50 µg/100 ml.

Group V Patients with haemoglobin under 10 g., normal MCHC, evidence of both hypochromic and macrocytic anaemia on the film and a low or low-normal serum iron, (all cases below 60 µg%).

Group VI Patients with a low haemoglobin, normal MCHC, evidence of macrocytic anaemia on the film, and a normal to high serum iron, (all cases had a value above 80 µg%).

RESULTS

(a) Haemoglobin distribution:

	0	10	20	30	40	50	60	Percent of total.
Less than 10 G %			20					11,4
10-10,9 G %				27				16,0
11-11,9 G %					49			28,1
12-12,9 G %						56		31,4
13-13,9 G %			20					11,4
14-14,9 G %	3							1,7

Mean haemoglobin = 11,6 G %
 11,4 % have values below
 10 G %

(b) Serum iron distribution:

	0	10	20	30	40	Percent of total
Less than 30		8				4,8
30-49			27			16,1
50-69				33		19,6
70-89					40	23,8
90-109			27			16,1
110-129		25				14,8
130-149	4					2,4
150+	4					2,4

Mean serum iron = 81 µg %
 20,9% have values below 50 µg %

(c) Average values

Parity	Haemoglobin
0	11,7
1	11,7
2	11,9
3	11,8
4	11,5
5	11,6
6	11,3
7	11,3
8	10,3
9	12,2
11	10,3
Total	11,6

(d) Classification

Group I (Normal)
Group II (Normal low)
Group III (Hypochromic iron)
Group IV (Hypochromic iron)
Group V (Mixed iron)
Group VI (Macrocytic iron)

The haemoglobin of patients (11,4 per cent) less than 10 grams per cent was favourably with iron is no worse than pregnant women was chosen to

(c) Average values by parity:—

Parity	Haemoglobin	MCHC	Serum Iron ($\mu\text{g}\%$)	No. in sample
0	11,7	30,0	84	44
1	11,7	30,0	80	27
2	11,9	30,7	71	20
3	11,8	30,4	77	17
4	11,5	30,1	86	16
5	11,6	30,0	73	16
6	11,3	30,0	76	19
7	11,3	30,5	86	11
8	10,3	30,3	70	3
9	12,2	32,1	53	1
11	10,3	27,8	112	1
Total	11,6	30,2	81	175

(d) Classification by groups:

	No. in group	per cent. of total
Group I (Normal values)	129	73,2
Group II (Normal but iron low)	26	15,5
Group III (Hypochromic, iron low)	5	3,0
Group IV (Hypochromic, iron normal)	6	3,3
Group V (Mixed anaemia, iron low)	6	3,3
Group VI (Macrocytic, iron normal)	3	1,7

DISCUSSION

The haemoglobin histogram shows that 20 patients (11,4 per cent.) had a haemoglobin value less than 10 grams per 100 ml. This compares favourably with Scott's findings in Glasgow, and is no worse than values found in other studies of pregnant women in Britain. The value of 10 G% was chosen to make an adequate allowance for

physiological haemodilution in pregnancy, and this value seems to be fairly widely accepted in obstetrics. Of the 20 anaemic patients 11 had hypochromic anaemia, 6 had mixed hypochromic and macrocytic, and 3 had pure macrocytic anaemia. Thus 17 (9,7 per cent.) had evidence of hypochromic anaemia and 9 (5,1 per cent.) some evidence of possible folic acid deficiency.

The serum iron histogram shows that 35 (20,9 per cent.) patients, or approximately one in every five, had a serum iron below 50 $\mu\text{g}\%$. The limits of normal in European patients are taken as 50 and 150 $\mu\text{g}\%$ ($100 \mu\text{g}\% \pm 2$ standard deviations). It is interesting to note that only 9 out of 35 patients with low serum iron values had significantly low haemoglobin values, although the average was lower than for the whole sample (10,9 G% — not statistically significant). 5 out of 8 patients with a serum iron of less than 30 $\mu\text{g}\%$, however, had a haemoglobin less than 10 G, with an average of 9,3 G (statistically significantly lower than the mean for all patients, 11,6 G%). Where serum iron values fell below 20 $\mu\text{g}\%$ all patients had a low haemoglobin value as well (average 7,8 G%). Thus while there is a general correlation between low serum iron and low haemoglobin values, it does not automatically follow that a patient with a low serum iron will be anaemic, or that a patient who is not anaemic has an adequate serum iron. This finding is consistent with the view that haemoglobin levels are maintained in preference to other iron stores in iron deficiency.

In the histories taken there was a remarkable lack of positive findings such as menorrhagia, abortions or ectopic pregnancy among those women found to be anaemic. Among those who had had such complications, the incidence of anaemia was no greater than that for the group as a whole. The abortion rate for all pregnancies in the women of the samples was 3,3 per cent., surprisingly lower than the expected 10-20 per cent. This may reflect a decreased tendency for those who abort to become pregnant again, or merely an increased interest in maintaining pregnancies in those women who attended this antenatal clinic.

In view of the high incidence of iron deficiency found in pregnant African women in this series, prophylactic iron therapy for every woman attending antenatal clinics would seem justified, regardless of the haemoglobin level found. Only four out of the series had a serum iron greater than 150 $\mu\text{g}\%$, so that the chances of iron overload do not seem to be very high. Since about 1 in 20 patients had either a mixed deficiency or a macrocytic anaemia, prophylactic folic acid

would appear to be indicated in those who are anaemic, especially as pernicious anaemia is very rare in African women of child bearing age. Serum folate determinations in a similar series would be most helpful in evaluating the true incidence of folic acid deficiency.

Contrary to what might have been expected, mean values for the haemoglobin, MCHC, and serum iron did not differ significantly on statistical analysis of the parity groups. Perhaps this is due to an average interval between pregnancies of 18 months to two years in the African multiparae in this series, or to a high dietary iron intake.

SUMMARY

(1) An account of an investigation into haemoglobin, MCHC and serum iron values in 175 African women attending Harare Hospital Antenatal Clinic for the first time in the current pregnancy is given.

(2) The mean haemoglobin for the group was 11,6 G%. 11,4% of the patients had values less

10 G%. 6,3 per cent. had evidence of hypochromic anaemia, 1,7 per cent macrocytic anaemia, and 3,3 per cent. had a mixed anaemia. The mean MCHC was 30,2 per cent.

(3) The mean serum iron was 81 μg per 100 ml. 20,9 per cent had levels below 50 μg %.

(4) There was no significant difference in haemoglobin, MCHC and serum iron values between groups of women of differing parity.

ACKNOWLEDGMENTS

I should like to thank Professor Philpott, Professor of the Department of Obstetrics and Gynaecology, University of Rhodesia, and Mr. A. Bedford, for their help to me in the planning and execution of the project.

REFERENCES

- DAVIDSON, L. A. G. (1964). *C. Afr. J. Med.* 10, 96. March, 1964.
HOWARD, J. K. (1966). *C. Afr. J. Med.* 12, 231. December, 1966.
DONALD, I. (1964). *Practical Obstetric problems*. Lloyd-Luke.



Toxicology of Animals and Plants

Toxins of Animal and Plant Origin: Vol. 2. Edited by A. de Vries and E. Kochva. Gordon & Breach, London, 1972, pp. 493-821. £8,10.

This is the second of three volumes which comprise the Proceedings of the Second International Symposium on Animal and Plant Toxins held at Tel Aviv, February, 1970. Volume 1 is divided into three sections: (a) General, (b) Venom glands and secretion of (c) Biochemistry (snakes). Volume 3 will contain sections on immunology and clinical aspects. Volume 1 was reviewed earlier.

This volume (Vol. 2) contains section III (11 papers) on Biochemistry (other species) and section IV (15 papers) on Pharmacology. 'Other species' refers to species other than snakes. All but two papers of the volume are in English. Section III describes various aspects of the isolation, purification, structure and chemistry of a diverse variety of venoms. The format and type

of information is analogous to that part of Volume 1 which was devoted to snake venoms.

The section on pharmacology presents information on the effects of snake and other venoms on neuromuscular and cardiovascular phenomena and blood coagulation. Properties and actions of toxins produced by various microorganisms are also presented.

No index is provided but no doubt a composite index will appear in Volume 3. However, a list of paper titles is given at the front of the book and this provides a great deal of the information required. Each paper is complete in itself and is followed by a list of references.

The volume provides a compendium of the latest definitive work on the biochemistry and pharmacology of venoms and toxins. This volume is of interest only to the basic research worker. Even though this book appeals only to the specialist worker, the price is phenomenally high.

D. P. THORNHILL.

is
● combats mi
by unidentifie
● combines t
tetracycline
coccal action

Albamy Capsules

Each capsule co
Novobiocin (as
Tetracycline hy
Supplied in bott

Albamy Flavoured Gr

When reconstitu
pension contain
Novobiocin (as
Tetracycline (e
hydrochloride
Potassium meta
Supplied in bott
(when reconstitu

728 REGISTERED TRADEM