

WORKING PAPER

No. 4-03

Awareness and the Demand for Environmental Quality:
Drinking Water in Urban India

Jyotsna Jalan
E. Somanathan
Saraswata Choudhuri



South Asian Network for Development
and Environmental Economics

December 2003

Awareness and the Demand for Environmental Quality: Drinking Water in Urban India

JYOTSNA JALAN
E. SOMANATHAN
SARASWATA CHOUDHURI

*Society for Economic Research and Financial Analysis
Indian Statistical Institute
New Delhi, India*

December 2003

South Asian Network for Development and Environmental Economics (SANDEE)
PO Box 8975, EPC 1056
Kathmandu, Nepal

Working Paper No. 4-03

Published by the South Asian Network for Development and Environmental Economics (SANDEE),
PO Box 8975 EPC 1056 Kathmandu, Nepal.
Telephone: 977-1-552 8761, 552 6391 Fax: 977-1-553 6786

SANDEE research reports are the output of research projects supported by the South Asian Network for Development and Environmental Economics. The reports have been peer reviewed and edited. A summary of the findings of SANDEE reports are also available as SANDEE Policy Briefs.

National Library of Nepal Catalogue Service:

Jyotsna Jalan

E.Somanathan

Saraswata Choudhuri

Awareness and the Demand for Environmental Quality: Drinking Water in Urban India

ISBN 99933-826-3-9

Key Words

1. Environmental Awareness
2. Drinking Water Quality
3. Health Risks
4. Willingness to Pay

The views expressed in this publication are those of the author and do not necessarily represent those of the South Asian Network for Development and Environmental Economics or its sponsors unless otherwise stated.

The South Asian Network for Development and Environmental Economics

The South Asian Network for Development and Environmental Economics (SANDEE) is a regional network that brings together analysts from different countries in South Asia to address environment-development problems. SANDEE's activities include research support, training, and information dissemination. SANDEE is supported by contributions from international donors and its members. Please see www.sandeeonline.org for further information about SANDEE.

Technical Editors:

Herath Gunatilake

Priya Shyamsundar

Comments should be sent to E. Somanathan, Indian Statistical Institute, 7 Shaheed Jit Singh Marg, New Delhi 110016, India. E-mail: som@isid.ac.in

TABLE OF CONTENTS

1. INTRODUCTION	1
2. DATA	3
3. BIVARIATE PROBIT MODEL	7
4. MULTINOMIAL LOGIT MODEL	10
5. EXPECTED WILLINGNESS TO PAY FOR POTABLE DRINKING WATER	14
6. CONCLUSION	20
7. ACKNOWLEDGEMENTS	20
REFERENCES	21
APPENDIX A : Construction of the wealth index	23
APPENDIX B : Estimated costs of different purification methods in city of Delhi	25

LIST OF TABLES

Table 1: Purification adoption rates for various household characteristics	5
Table 2: Marginal effects for the bivariate probit model	8
Table 3: Marginal effects of the multinomial logit model	12
Table 4: Descriptive statistics for expected willingness-to-pay for Delhi (Lower bound in Indian rupees)	15

LIST OF FIGURES

Figure 1: Kernel densities associated with estimated willingness to pay for the different wealth quartiles	17
Figure 2: Kernel densities associated with estimated willingness to pay for the different types of media exposure	17
Figure 3: Kernel densities associated with estimated willingness to pay for the different levels of adult female education	18
Figure 4: Kernel densities associated with estimated willingness to pay for the different levels of adult male education	18
Figure 5: Wealth and female education effects on WTP	19
Figure 6: Wealth and male education effects on WTP	19

Abstract

The demand for environmental quality clean air, potable water, sanitation, safe food is often presumed to be low in developing countries due to poverty. However, individuals in developing countries often lack the necessary information to make good decisions about environmental hazards in their day-to-day lives. Even if households can afford to take private measures to improve environmental quality, very often they choose not to do so, because they are not aware of the health risks associated with inferior environmental quality. A key policy question is whether increasing awareness about the adverse health effects of environmental pollution will increase demand for a cleaner environment?

In this paper, a household survey from urban India is used to estimate the effects of awareness and wealth on household decisions to purify home water. Average costs of different home purification methods are used to get estimates on willingness to pay for better drinking water quality in Delhi. It is found that measures of awareness such as schooling and exposure to mass media have statistically significant effects on adoption of different home purification methods and therefore, on willingness to pay. The interesting result is that these effects are similar in magnitude to wealth effects — this suggests that lack of awareness may be as important as poverty in influencing demand for clean water.

JEL Classification Numbers: O13, Q25

Keywords: Environmental awareness, drinking water quality, health risks, willingness to pay.

Awareness and the Demand for Environmental Quality: Drinking Water in Urban India

Jyotsna Jalan, E. Somanathan and Saraswata Choudhuri

1. Introduction

The demand for environmental quality - clean air, potable water, sanitation, safe food - is often presumed to be low in developing countries due to poverty and this is considered to be efficient since it reflects choices made by individuals in their own best interests. However, individuals in developing countries often lack the necessary information to make good decisions about environmental hazards in their day-to-day lives.

In this paper, we show that even if households can afford to take private measures to improve their environmental quality, very often they choose not to do so, because they are not aware of the health risks associated with inferior environmental quality.

Contaminated drinking water is a major health hazard in developing countries and diarrhoea is the most common disease associated with it. This is particularly true for the poor and the vulnerable. The World Health Organization (WHO, 2002) estimates 1.7 million deaths and 54.2 million disability adjusted life years (DALYs) lost worldwide per year due to unsafe water, hygiene and sanitation.¹ Almost all of these deaths are in developing countries and nine out of ten deaths occur in children. In India alone, there are more than a million child deaths per year resulting from waterborne diseases like diarrhoea (Parikh et. al., 1999).

Interventions to improve drinking water quality and minimize exposure to health risks from ingestion of contaminated water include provision of quality regulated piped water supply to all households. According to a WHO (2002) estimate, if universal piped and regulated water supply were achieved, 7.6 billion diarrhoea cases could be averted annually worldwide. However, such state intervention is unlikely to be forthcoming in the short-run, especially in the absence of a clearly expressed demand from the public, which in turn depends on public awareness of the health hazards of drinking contaminated water.

A substitute short-term solution is dis-infection at point of use. This includes low-grade technologies like straining with a cloth, using chlorine and safe storage vessels, to relatively more sophisticated and expensive technologies like electronic filters that use ultra-violet radiation to remove pathogens etc. These interventions have large health impacts in regions of high child mortality and some are inexpensive. Straining with a cloth for example, especially if it is folded over to form eight or more layers, has been shown to be successful in removing copepods, which are a host for cholera bacteria (Colwell et. al., 2003).

The question is why has adoption of safe drinking water practices not been universal? Poverty is an important factor but it certainly cannot explain lack of use of even costless methods like straining with a cloth. According to a study in rural Bolivia, cited in Gadgil (1998), many people were unaware of the link between water contamination and diarrhoea. Undoubtedly, awareness is a determinant of the demand for safe drinking water. In this paper we estimate the magnitude of the “awareness” effect and compare it to that of wealth.

¹ One DALY being equal to the loss of one healthy life year.

We use measures of schooling, exposure to mass media, and occurrence of diarrhoeal disease as indicators of a household's awareness about the health risks associated with drinking contaminated water. The first two are self-explanatory. Regarding the third, it is possible that a household afflicted with waterborne diseases in the past will take some steps to improve its quality of drinking water to reduce the likelihood of such disease in the future.

While we are not aware of any previous research specifically examining the effect of awareness on water purification in developing countries, studies on demand for water have found evidence that education is one of the determinants. Dasgupta (2001) and McConnell and Rosado (2000) use data from Delhi and an urban area in Brazil respectively to show that education of the household head is statistically significant in a household's decision to purify its drinking water controlling for income and other household characteristics.^{2,3} When formulating public health policy however, it is important to distinguish whether information is gathered through formal schooling, the mass media, or knowledge gained through experience of a previous health shock. This is what we do in this paper - we examine the role of different sources of knowledge in adoption of environmental risk averting behavior by households.

We summarize the main findings of our paper here. Each of the awareness indicators included in the analysis has an independent (of household wealth effects) significant effect on a household's decision to adopt some home purification method. The magnitudes of these effects are by no means small and are comparable to the wealth effects. For example, a one-year increase in the years of schooling of the most educated female member of the household increases the probability of home purification by about 1.4 percentage points. Since this variable ranges from 0 to 23 years, its effect is comparable in magnitude to the increase of 24.6 percentage points resulting from a move from the first to the fourth wealth quartile. The probability of purification rises by 8 percentage points if a female household member reads a newspaper at least once a week. A prior bad health experience raises the probability that households boil their drinking water by about 5 percentage points.

Finally, using estimated probabilities of the different methods of purification from the econometric model, and average costs of the different methods from an informal survey of markets in the metropolitan city of Delhi, we calculate the expected cost of improving household water quality for the sampled households in Delhi. Our estimates indicate that expected willingness to pay increases by nearly 50 percent from Rs. 212 per person per year to Rs. 314 when the schooling of the most educated female in the household rises from 3.5 to 10.5 years. By comparison the expected willingness to pay rises from Rs. 212 to Rs. 289 when moving from the second to the third wealth quartile. It rises by over 25 percent when a woman member of the household reads a newspaper at least once a week.

The following section describes the data and reports summary statistics. Sections 3 and 4 present the results from bivariate probit and multinomial logit models respectively that are used to estimate the role of awareness in the household's decision-making process for home purification of drinking water. Section 5 estimates willingness to pay for households in the city of Delhi. Section 6 concludes.

² Alberini *et. al.*'s (1996) study of diarrhoea and avoidance behavior through hand-washing in Jakarta, Indonesia, does not show any significant effect of mother's education on either diarrhoea or avoidance behavior. In their study, boiling drinking water was pervasive, reflecting its low cost owing to the fuel subsidies in Jakarta. This is very far from being the case in urban India. Another related study is the one by Larson and Gnedenko (1999) of home water purification in Moscow, in which, again the cost of boiling is low, and protection from pathogens does not appear to be the main concern.

³ There are also plant or community-level studies of industrial pollution such as that of Pargal and Wheeler (1996) who relate the level of schooling in a community and other variables to the pollution level from plants in the neighborhood. In one of their specifications, schooling significantly reduces pollution, perhaps reflecting collective action that pressures the plant to lower pollution. Evidence from the US, also at the community level (Brooks and Sethi, 1997) or at the state level (Boyce *et. al.*, 1999) indicates that schooling seems to lower pollution levels.

2. Data

We use data from the National Family Health Survey of India, (NFHS), conducted by the International Institute for Population Sciences in 1998-1999. The NFHS sample covers 99 percent of India's total population living in 25 out of 26 states (Tripura and union territories were not included) and is a household survey with an overall target sample size of approximately 90,000 ever-married women in the age group 15-49.⁴

The sample households are evidently highly exposed to waterborne disease. One of the survey questions asked whether there were any cases of diarrhoea among children aged 0-3 years in the two weeks preceding the survey. Of the 7,351 households that had children in this age group, 1,014 or 14 percent reported at least one case in the preceding two weeks.

Among the households who treat their drinking water, four different methods are used– (i) straining water with an ordinary cloth, (ii) using alum (aluminum sulphate, a flocculant) tablets, (iii) using an ordinary (candle) filter, (iv) using an electronic filter or (v) boiling water. The choices are mentioned in ascending order of their cost.^{5,6}

As mentioned earlier, straining with a cloth folded over to form at least 8 layers can provide significant protection against cholera, though most free-floating bacteria would pass through the mesh (Colwell *et. al.*, 2003).⁷

Water can also be treated with flocculants like potassium and aluminum sulfates. Flocculants cause small particles including bacteria to form clumps that can be allowed to settle or that can be filtered using ordinary filters. These filters are also effective in reducing turbidity and in large reductions in bacterial and other pathogens like giardia, amoebae, etc. However, ordinary filters do not themselves disinfect water adequately i.e. will not remove all fecal pathogens.

Electronic filters first filter particulate matter and then irradiate the water with ultraviolet light, killing all pathogens. Provided they are maintained well, electronic filters are as effective as boiling at removing pathogens. Electronic filters impart no taste or odor to the water, and consume far less energy than boiling water does.

Boiling is the oldest method to obtain drinking water free of biological contaminants. In many developing countries, residents routinely boil their water because they perceive their water supply to be unsafe for drinking purposes. The World Bank estimates that 1 percent of Jakarta's GDP is spent on boiling drinking water.

In our analysis, for households using more than one method of purification, we have considered only the 'best' method (in terms of removal of pathogens from the water) and we have clubbed together households using alum tablets or ordinary filters because there were very few households using the

⁴ We included only urban households with 15 or fewer members. Furthermore, we exclude households for which some variables were missing. Accordingly we retain 20,681 of the 25,243 households in the original sample.

⁵ We do not consider treatment to remove chemicals from drinking water. While chemical contamination is important in certain localities, in poor countries like India, the main adverse health effects from water contamination are due to biological contamination from pathogens that originate from fecal matter (Gadgil, 1998).

⁶ Costs are calculated in Appendix A.

⁷ In the same study, a large number of mothers using cloth purification reported a decline in the incidence of diarrhoea within families.

former in our sample.⁸ Furthermore, given that the NFHS does not report the percentage of total drinking water consumption that a household purifies, we assume that a household's entire drinking water consumption is treated.

Table 1 reports purification adoption rates for households with different characteristics. Despite the high exposure to diarrhoeal disease, a significant 47 percent of households continue to consume water without using any type of water purification method. This percentage remains as high as 32 percent even for households in the top wealth quartile.⁹ Furthermore, only about 20 percent of households use methods (i.e. boiling and electronic filters) that reliably remove commonly found biological pathogens.¹⁰

There is no direct information in the NFHS about how informed a household is regarding the health risks arising from drinking contaminated water. But our hypothesis is that if the adult members of a household, who are presumably the decision makers, are educated, it is likely that they will be aware of the health risks associated with drinking non-potable water. We therefore use education of adult members to proxy for health awareness of households. The NFHS collected information on the educational qualifications of individuals. We use highest number of years of education among female household members aged 15+, and highest number of years of education among male household members aged 15+ to capture education levels of adult male and female members in the household.

Table 1 shows that the proportion of households who do not purify their water is smaller for households with higher adult human capital stocks. However, even among households whose adult members are highly educated (i.e. have 14+ years of education), 25-30 percent do not use any water purification method and even among those who adopt some method, about 40 percent use methods that are not fully effective in removing biological pathogens from the water.

The NFHS does not collect information on consumption or income. However, the survey collects information on household ownership of various assets (car, motorcycle, bicycle, refrigerator, telephone, color television, black and white television, radio, mattress, clock or watch, table, chair, etc.) and characteristics of household dwelling (house type, toilet type, number of rooms per capita, source of light, fuel type, utensils type, etc.). We use the first principal component of these variables as a wealth index. Appendix A provides the formula for the wealth index. Filmer and Pritchett (2001) used an almost identical index from the first round of the NFHS and found that it explained consumer expenditure data quite well.

The first principal component explains 24.36 percent of the variation in the asset variables. For ease of interpretation, we create and use wealth quartiles from the wealth index rather than the actual wealth index in our analysis.¹¹ Households in the bottom quartile correspond to the poorer units in the sample.

One concern is that our wealth index is not comprehensive so the information/awareness variables could still act as a proxy for unobserved wealth effects. To examine the plausibility of this speculation, we created another index with all the variables used earlier together with the education variables mentioned above. Spearman's rank correlation coefficient ($r=0.9878$) between the two indices shows that the ranking of the households changes very little with or without the education variables in the wealth index.

⁸ We have arbitrarily ranked electronic filtration to be more effective than boiling. Since very few households engage in both, this particular choice is unlikely to make any difference to the results.

⁹ We explain below how data on wealth are constructed.

¹⁰ Less than 3 percent of households who use some purification method use an electronic filter. It is only in the last few years that there has been a proliferation of such products both in terms of availability in the market and usage by consumers.

¹¹ We did estimate a bivariate probit model with the wealth index rather than its quartiles. The coefficients on the awareness variables changed very little.

Table 1: Purification adoption rates for various household characteristics

	No purification	Purification			
		Straining	Alum/ordinary filter	Electronic filter	Boiling
<i>Wealth:</i>					
Bottom quartile	66.16	19.48	2.64	0.08	11.64
Second quartile	50.16	22.32	7.08	0.24	20.16
Third quartile	41.56	20.84	15.92	0.48	21.20
Top quartile	31.96	11.36	28.96	5.12	22.60
<i>Highest number of years of education among adult female household members:</i>					
No education	68.84	20.67	2.71	0.10	7.67
1-7 years	54.98	23.35	6.69	0.23	14.74
8-13 years	42.57	17.87	15.36	1.07	23.13
14+ years	26.32	11.68	30.31	5.49	26.20
<i>Highest number of years of education among adult male household members:</i>					
No education	67.67	16.37	4.90	0.31	10.74
1-7 years	59.86	22.30	4.51	0.12	13.22
8-13 years	48.19	20.20	10.83	0.68	20.10
14+ years	30.19	13.57	28.48	4.36	23.40
<i>Exposure to mass media:</i>					
Doesn't read newspaper once a week	60.45	19.96	6.37	0.37	12.85
Reads newspaper at least once a week	32.12	16.79	22.26	2.79	26.04
Doesn't watch TV once a week	62.73	19.81	3.80	0.19	13.48
Watches TV at least once a week	44.08	18.22	15.83	1.77	20.10
Does not listen to radio once a week	53.16	21.15	10.53	1.10	14.07
Listens to radio at least once a week	41.55	15.76	16.90	1.87	23.91
All households	47.45	18.50	13.66	1.48	18.90

Note: The sample consists of 20,681 households from urban India in 1998-99.

Simple descriptive statistics suggest important wealth effects in the adoption of water purification methods. Households belonging to the upper wealth quartiles used costly purification methods like an ordinary water filter or alum, electronic filter, or boiling. Thus, while 2.64 percent of households in the bottom quartile used alum tablets or an ordinary filter, this number increases to 30 percent in the uppermost quartile. Nearly two-thirds of households in the bottom wealth quartile used no water purification method; in the fourth quartile, this proportion was less than a third. Electronic filters were primarily used by households in the top wealth quartile. (See Table 1 for further details).

Mass media in India often report on episodes of waterborne diseases and precautions that can be taken especially during the rainy season when incidence of such diseases increase. Government agencies sponsor social awareness messages relating to prevention of waterborne diseases in the media. Since the NFHS collects data on women's exposure to such mass media, we include variables like whether any female member of the household reads a newspaper, watches television, or listens to the radio at least once a week in our specifications.¹²

Among households not exposed to the print media, sixty percent do not treat their water. This proportion falls by half for households that read newspapers. This differential was somewhat less stark for two of the other media variables: watching TV and/or listening to the radio at least once a week.

Lastly, it is plausible that a household's perception about the health risks from drinking unsafe water is influenced by a health "shock" in the past that can be related to poor drinking water quality. That is, households learn from a bad health experience and their willingness to adopt safe drinking water practices increases. We use occurrence of diarrhoea in the two weeks preceding the survey, in the age group 0-3 years to measure household's exposure to waterborne diseases. This variable incorporates some information, albeit quite limited, on household morbidity due to drinking water quality.

We also include other control variables: state dummies, demographic characteristics (household size, household head's age, age squared, and marital status, proportion of household members in age groups 0-3 years, 3-5 years, and 5-15 years, socio-economic variables (religion and caste dummies, proportion of household members in different occupations (those not working, in occupations that (do not) require higher education, in medical professions, and in the service sector)), place of treatment when ill and source of water dummies, (whether piped into home, a public tap, or other source such as a handpump or borewell).^{13,14}

Descriptive results reported in this section suggest strong unconditional correlations between household's demand for improved quality of drinking water and the different awareness indicators. However, many of these variables are highly correlated with each other and the observed patterns based on estimated unconditional correlations could be misleading. We thus need to develop an econometric model where we are able to separate the individual effects of each of the above variables from the others. We do this in the next two sections.

¹² We also include whether any female member of the household goes to the cinema at least once a month in our analysis but this variable was not significant in practice.

¹³ As some of the states few observations, we clustered them with other states based on geographic, socio-economic and cultural proximity.

¹⁴ Jalan and Ravallion (2002) find that provision of piped water to a household is not adequate to prevent diarrhoea among children. Prevention rates are higher if the infrastructure (piped water) is combined with the stock of human capital of household members particularly those of the female members.

3. Bivariate Probit Model

A household's perception of the health risks from drinking water and therefore its purification behavior, could be affected by its past experience of water-borne disease. Of course, its experience of diarrhoea will be conditional on its purification behavior. To allow for this, we begin by estimating a bi-variate probit model of purification and occurrence of diarrhoea among 0-3 year olds in the household.¹⁵

We have data on incidents of diarrhoea among 0-3 year olds in the two weeks preceding the day of survey that we use as a proxy for prior incidence of diarrhoea. We want to assess whether a household's decision to purify water by any means depends on its information/awareness about the advantages of improving the quality of water, independently of the household's wealth status.

Our estimated econometric specification is:

$$\begin{aligned}
 y_1^* &= \mathbf{b}'_1 x_1 + \mathbf{e}_1, \quad y_1 = 1 \quad \text{if} \quad y_1^* > 0, 0 \text{ otherwise} \\
 y_2^* &= \mathbf{b}'_2 x_2 + \mathbf{e}_2, \quad y_2 = 1 \quad \text{if} \quad y_2^* > 0, 0 \text{ otherwise} \\
 E(\mathbf{e}_1) &= E(\mathbf{e}_2) = 0 \\
 \text{Var}(\mathbf{e}_1) &= \text{Var}(\mathbf{e}_2) = \mathbf{s}^2 \\
 \text{Cov}(\mathbf{e}_1, \mathbf{e}_2) &= \mathbf{r}
 \end{aligned} \tag{1}$$

The binary variables y_1 and y_2 are indicators for whether or not a household adopts any home purification of water, and whether or not a 0-3 year old in the household has been afflicted by diarrhoea in the two weeks preceding the survey. x_i 's are the vector of explanatory variables (described in Section 2 above) that includes the wealth, education and media exposure variables and \mathbf{e}_i 's are correlated, identically distributed random variables. We estimate this model using maximum likelihood techniques. The correlation coefficient between the errors of the two equations is statistically significant (the Likelihood Ratio Test for $H_0: \rho=0$ against $H_1: \rho \neq 0$ gave a p-value of 0.0329) thus rejecting the hypothesis that the two dependent variables are not jointly determined. The results are reported in Table 2.¹⁶

The probability of purification is 7 percent greater for households in the second as compared to the first wealth quartile, and as much as 24.5 percent greater for those in the fourth as compared to the first quartile. Educational attainments of adult household – male or female – members are also important in the household's decision to purify water. A single additional year of education among the highest educated female (male) adults in the household will increase the probability of using some method of purification by 1.4 (1.0) percentage points. Two of the media dummies: whether any female household member reads a newspaper at least once a week, and whether she listens to the radio at least once week, are statistically significant. It should be noted that even though possession of a television or radio are components of the wealth index, they are included independently in the specification so that watching TV and listening to the radio do not proxy for ownership of these media.

¹⁵ Ideally we would have liked to include the incidence of diarrhoea among all household members but this information is not available in the data.

¹⁶ Household water purification and occurrence of diarrhoea are assumed to be jointly determined in this estimation procedure.

Table 2: Marginal effects for the bivariate probit model

	Probability of purification	Probability of occurrence of diarrhoea
Wealth quartile 2	0.0724* (0.0124)	-0.0033* (0.0014)
Wealth quartile 3	0.1255* (0.0147)	-0.0037* (0.0017)
Wealth quartile 4	0.2458* (0.0167)	-0.0058* (0.0022)
Maximum years of education among female members	0.0143* (0.0012)	-0.0000 (0.0002)
Maximum years of education among male members	0.0096* (0.0014)	-0.0001 (0.0001)
Female members read newspaper at least once a week	0.0807* (0.0122)	-0.0003 (0.0015)
Female members watch television at least once a week	0.0051 (0.0125)	0.0025 (0.0016)
Female members listen to the radio at least once a week	0.0340* (0.0096)	0.0003 (0.0013)
Number of observations	20,681	
Log likelihood	13,386.38	

Standard errors reported in parentheses.

* indicates significance at 5 percent or lower.

A different specification in which the wealth index was included directly in place of quartile dummies yielded very similar coefficients and marginal effects for the awareness variables. We also estimated a model that included interactions between the wealth quartiles and other variables, but nearly all the interaction terms were statistically insignificant and the coefficients of interest were very similar.

The second column of Table 2 reports the marginal effects on the probability of the occurrence of diarrhoea. Only wealth is statistically significant.

4. Multinomial Logit Model

To study the effects of awareness on willingness to pay for water quality, we need to allow for the fact that different purification methods have different costs. To examine this, we estimate a multinomial logit model where a household has five different choices available – no purification, straining with a cloth, alum tablets or ordinary (candle) water filter, electronic filtration, or boiling.

Let U_{ij} be the i^{th} household's utility from making the j^{th} purification choice for $j = 0,1,2,3,4$ where $j = 0$ denotes the choice of not purifying. The observed choice of j^{th} purification method used by the i^{th} household is defined as:

$$Y_{ij} = 1 \quad \text{if } Y_{ij}^* = \text{Max}(Y_{i0}^*, Y_{i1}^*, \dots, Y_{i4}^*) \quad (2)$$

$$= 0 \text{ otherwise}$$

The intuition underlying this model is that a household faced with many alternatives makes a single decision depending on which alternative gives it the highest utility.

The underlying latent variable is specified as:

$$Y_{ij}^* = \beta_j x_i + \varepsilon_{ij} \quad (3)$$

where x_i is the vector of observed household characteristics, and ε_{ij} is a residual that captures unobserved household characteristics and errors in optimization by the household. The ε_{ij} 's are assumed to be identically and independently distributed with Type I extreme-value distribution. It is assumed that the household chooses only one type of purification method.¹⁷ Standard numerical optimization algorithms can be used to estimate the model.

Finally, we have to allow for the endogeneity of occurrence of diarrhoea. Given that estimating a simultaneous equation multinomial model is computationally very burdensome, we use a two-stage procedure. In the first stage a logit model of the probability of getting diarrhoea among 0-3 year olds as a function of the explanatory variables in the previous model as well as the following additional variables: knowledge of adult women regarding oral re-hydration (ORS) products (whether a woman has heard of ORS before, recognizes the ORS packet, used ORS), recognition of symptoms of diarrhoea by women (repeated watery stools and vomiting, blood in stools etc.), is estimated. In the second stage, the multinomial logit model is estimated in which the predicted probability from the logit model is included as a regressor in addition to the explanatory variables from the previous model.

The estimated coefficients in the multinomial model are difficult to interpret. We need to calculate the marginal effects for the variables of interest (i.e. the change in the probability of adoption of a particular alternative with a unit change in the explanatory variables) which are not only functions of their own estimated coefficients but also depend on all other estimated coefficients and levels of regressors.

¹⁷ In the data however, some households may have indicated adoption of more than one purification method. In order to cut down the number of alternatives we have ranked the different purification methods and assumed that the household adopts the most effective (in terms of removing pathogens) method.

The analytical expressions for the marginal effects are for any $m=0,\dots,4$:

$$\frac{\partial P(Y = m | X)}{\partial x_k} = P(Y = m | X) \cdot [\beta_{mk} - \sum_{j=0}^4 \beta_{jk} P(Y = j | X)] \text{ , if } x_k \text{ is continuous} \quad (4)$$

$$\frac{\Delta P(Y = m | X)}{\Delta x_k} = P(Y = m | X, x_k = 1) - P(Y = m | X, x_k = 0) \text{ , if } x_k \text{ is discrete}^{18} \quad (5)$$

The response probabilities are given by:

$$P(Y = m) = \frac{e^{\beta_m x}}{1 + \sum_{j=1}^4 e^{\beta_j x}} \text{ for } m = 1, 2, 3, 4 \quad (6)$$

$$P(Y = 0) = \frac{1}{1 + \sum_{j=1}^4 e^{\beta_j x}} \quad (7)$$

In expressions (4) and (5), β_m is the coefficient associated with the m^{th} choice in the multinomial logit model.

The analytical standard errors of the marginal effects for our sample (20,681 observations) and model (more than 20 explanatory variables) are computationally very burdensome to calculate. Hence, we bootstrap our multinomial model to estimate the associated standard errors. We use a sample size of 20,000 observations and 50 repetitions in our computations.

We now turn to the results from the multinomial logit model. In the first stage logit equation of occurrence of diarrhoea among 0-3 year olds, the variables serving as excluded regressors from the second stage, knowledge of the household members regarding oral rehydration products (whether member has heard of ORS before, recognizes the ORS packet etc.), recognition of symptoms of diarrhoea by household members (repeated watery stools and vomiting, blood in stools etc.) are highly statistically significant. The estimated pseudo R^2 is very high for cross-sectional data at 0.52. Using this model we compute the estimated probability of occurrence of diarrhoea that we use in the second-stage multinomial model of the different purification methods.

Our estimates of the multinomial logit model are reported in Table 3. The first column reports the estimated marginal effects of the different variables on the probability of no purification evaluated at the means of the explanatory variables. In congruence with the bivariate probit model, relative to the bottom-most wealth quartile, households in the other three quartiles are more likely to adopt some kind of purification method to improve their quality of drinking water.

In the bivariate probit model, our unconditional marginal effects for purification showed a strong relationship between whether a household adopts a water purification method and both the wealth position and the “knowledge stock” of the household. Is this pattern also observed across different purification methods?

¹⁸ In case of categorical explanatory variables we report the change in probability of adoption of a particular alternative with a change from one state to another.

Table 3: Marginal effects of the multinomial logit model

	No purification	Straining	Alum or ordinary water filter	Electronic filter	Boiling
Wealth quartile 2	-0.1113* (0.0162)	0.0120 (0.0077)	0.0581* (0.0260)	0.0025 (0.1318)	0.0387* (0.0151)
Wealth quartile 3	-0.1965* (0.0181)	0.0047 (0.0077)	0.1409* (0.0361)	0.0025 (0.1337)	0.0485* (0.0174)
Wealth quartile 4	-0.3052* (0.0248)	-0.0392* (0.0099)	0.2488* (0.0521)	0.0207 (0.1374)	0.0750* (0.0207)
Maximum years of education among female members	-0.0161* (0.0015)	-0.0002 (0.0006)	0.0061* (0.0008)	0.0005* (0.0002)	0.0097* (0.0011)
Maximum years of education among male members	-0.0100* (0.0014)	0.0006 (0.0006)	0.0050* (0.0007)	0.0004* (0.0001)	0.0041* (0.0010)
Female members read newspaper at least once a week	-0.0776* (0.0119)	0.0049 (0.0068)	0.0291* (0.0051)	0.0011 (0.0007)	0.0425* (0.0089)
Female members watches television at least once a week	-0.0104 (0.0153)	0.0094 (0.0062)	0.0032 (0.0093)	0.0005 (0.0010)	-0.0027 (0.0131)
Female members listens to the radio at least once a week	-0.0401* (0.0085)	0.0142* (0.0055)	-0.0044 (0.0052)	-0.0003 (0.0004)	0.0306* (0.0084)
Children 0-3 years afflicted with diarrhoea in last two weeks at time of survey (predicted)	-0.0263 (0.0308)	0.0124 (0.0135)	-0.0365 (0.0288)	0.0038* (0.0017)	0.0467* (0.0239)
Observed frequencies	9,814	3,827	2,825	306	3,909
Number of observations	20,681				
Log likelihood	-15,830.89				

Standard errors reported in parentheses.

* indicates significance at 5 percent or lower.

Use of a costless but inferior purification method like straining with a cloth is less likely in the top quartile of the wealth distribution. Wealth has a large effect on the household's decision to adopt a purification method that entails some cost to the household. For example, relative to the bottom wealth quartile, the probability of an average household in the second quartile using a candle filter/alum to purify its water is 6 percentage points greater, and the same difference is 25 percentage points for the upper most wealth quartile. For the most expensive and common method of purification – boiling – we observe a similar trend where the marginal effects rise as we move up the wealth quartiles although not by as much.

We also want to test whether controlling for wealth, knowledge acquired by adult members through formal education leads to the household adopting superior methods of purification. Table 3 indicates that our education variables are significant with the education of the female adult members being the dominant effect. These effects are large: a *single* additional year of female education increases the probability of boiling by 1 percentage point compared to an effect of 0.47 due to an increase of one year of education among male adult members.

Among the media exposure variables used as proxies to capture transmission of knowledge to households through targeted information, listening to the radio and reading the newspaper are significant. It is interesting that for a cost-free purification method like straining with a cloth, female members listening to the radio is likely to increase the probability by 1.4 percentage points while for boiling, the probability increases by 3 percentage points.

The health experience proxy that we have used in our analysis - occurrence of diarrhoea among 0-3 year olds - is not significant for any of the purification methods except for boiling and electronic filter. Again this result is not unexpected since these two methods are the only ones that kill all bacteria.

Finally, in addition to the above variables, as mentioned in the data section we also included several control variables in our model. While details of their effects on the different home purification methods are available from the authors, it is worthwhile to comment on the effects of the occupation dummies. Model estimates suggest that households where the head works in the medical sector (surgeons, physicians, nurses etc.) are more likely to use alum tablets and or the ordinary filter compared to the others. However, there are no statistically significant effects of working in the medical sector on adoption of other purification methods.

5. Expected willingness to pay for potable drinking water

From a policy perspective, it is important to get an estimate of willingness to pay (WTP) for safe drinking water. Is it the case that better informed people are willing to pay more for a better quality environmental commodity than the uninformed? However, the NFHS was not designed as a contingent valuation survey nor does it collect information that will allow us to get an estimate of the household's willingness to pay. We can however use the probabilities predicted by the multinomial model and our calculated costs for the different methods of purification to estimate the lower bound that each household is willing to pay for safe drinking water.

We collected cost information on different brands of electronic filters, ordinary filters and alum tablets from a survey of local markets in the metropolitan city of Delhi.¹⁹ Under certain assumptions and based on the average cost price of these technologies, we estimated an annual per person cost of purification. The details on the assumptions made are spelt out in *Appendix B* of the paper. According to our estimates boiling is the most expensive method of improving the quality of drinking water. On the other hand, even though our data shows that use of electronic filter is not common in our sample, it is less than half as expensive as boiling and if maintained properly, improves the quality of water to the same extent as boiling does.

Let C_{ij} be the actual cost associated with the outcome j for household i . We assume that $C_{ij} = C_j \forall i$ i.e., cost of purification method j is constant over the entire population for each of the four purification methods. Under this assumption, C_j at best can be interpreted as the lower bound for willingness to pay by households for purification method j . The expected lower bound of willingness to pay (EW_i) by i^{th} household is given by:

$$EW_i = C_0 * P(Y_i = 0) + C_1 * P(Y_i = 1) + C_2 * P(Y_i = 2) + C_3 * P(Y_i = 3) + C_4 * P(Y_i = 4) \quad (8)$$

$P(Y_i=j)$ for $j=0,1,2,3,4$ is the associated estimated probability of adoption of purification method j for the i^{th} household from the multinomial model. We assume that all decision makers within the household will have identical $P(Y_i=j)$'s.

Using the sub-sample of households from Delhi only, estimated costs and probabilities of adoption of different purification methods, we estimate the (lower bound of) per person mean willingness to pay to improve the quality of water as Rs. 303 and per person median willingness to pay is Rs. 304 per year. (\$US 1 = Rs. 46)

How do these willingness to pay estimates relate to household wealth, education and media exposure variables? We report two sets of statistics. We fix values of all variables other than the variable of interest at their mean levels and report the expected willingness to pay. We thereby isolate the effects of the specific variable of interest from other household characteristics. We call this the "controlled" WTP. This is reported in the first column in Table 4. We also report the mean willingness to pay for different categories without controlling for household characteristics. For this latter case we report the mean, median and the standard deviation.

The controlled willingness to pay more than doubles from Rs. 158 in the bottom wealth quartile to Rs. 329 in the top quartile. Median willingness to pay increases ten-fold between the bottom and top wealth quartiles from Rs. 44 to Rs. 451.

¹⁹ We thank Dwiraj Bose for conducting the survey.

Table 4: Descriptive statistics for expected willingness-to-pay for Delhi (lower bound in Indian rupees)

	Controlled WTP	Mean WTP	Median WTP	Standard deviation
Wealth:				
Bottom quartile	158	59	44	56
Second quartile	218	136	110	86
Third quartile	252	235	214	119
Top quartile	329	434	451	124
Highest number of years of education among adult female household members:*				
No education	171	91	81	61
1-7 years	212	178	155	108
8-13 years	314	343	351	132
14+ years	426	497	501	93
Highest number of years of education among adult male household members:*				
No education	210	108	71	114
1-7 years	232	139	111	104
8-13 years	281	277	270	151
14+ years	328	462	480	121
Exposure to mass media:				
Does not read newspaper at least once a week	243	174	144	119
Reads newspaper at least once a week	309	429	451	130
Does not listen to radio at least once a week	57	226	195	154
Listens to radio at least once a week	65	369	402	171
Does not watch TV at least once a week	58	138	97	115
Watches TV at least once a week	61	320	327	175

Controlled WTP: Willingness to pay of the different sub-samples (e.g. bottom wealth quartile) keeping all other variables in the at their mean levels

Mean and median WTP: Respective sample descriptive statistics of the expected WTP

*For the two education variables, the controlled WTP is evaluated at 0, 3.5, 10.5 and 17 years of education to represent the four categories (no education, 1-7, 8-13 and 14+ years of education) respectively

The adult education variables have comparable effects. Median willingness to pay in the lowest category for maximum female education, no education, is Rs. 81, while for the highest category of 14+ years of education, it is Rs 497. The increase in median willingness to pay between no education and 1-7 years of education and no education and 8-13 years of education is greater for female than male education.

The controlled WTPs show a similar trend²⁰. The controlled WTP is lower in households where female members are uneducated compared to households where male members are illiterate. Furthermore the slope for female education years is steeper than that for education among males. For example, when number of years of education among female members increases from 3.5 to 10.5 years, WTP increases by 50 percent. However, for males the corresponding number is 21 percent. These numbers would probably support a view existing in the literature that men tend to judge health risks to be much smaller and less problematic than women do (WHO 2002, page 35).

Reading a newspaper is the most influential mass medium in terms of changing individual health risk perceptions associated with drinking unsafe water. The median willingness to pay is Rs. 451 and is comparable to the expected willingness to pay of the wealthiest and the most educated in the sample. This is also considerably higher when compared to the willingness to pay-Rs. 144-for improved quality of drinking water by the sample of households where female members do not read a newspaper even once a week. The difference between willingness to pay for a better quality environmental good among households exposed to the other two mass media-watching TV and listening to the radio once a week-compared to households not exposed to such mass media is equally significant.

The controlled WTPs for the media variables show a large difference between households where female members read a newspaper and where they do not-WTP is nearly 40 percent higher in households where female members read a newspaper at least once a week! The impacts of the other two media variables on WTP are less significant. Table 4 reports the details on these statistics.

Figures 1-4 depict the kernel densities associated with estimated willingness to pay for the different wealth quartiles, for different years of education across genders and for exposure to mass media. The pictures depict very clearly that expected willingness to pay for improved quality of drinking water rises with the wealth quartiles, with more education and with greater exposure to media variables.

²⁰ For both education variables, the controlled WTP's are evaluated at no education, at 3.5, 10.5 and at 17 years of education respectively.

Figure 1: Kernel densities associated with estimated willingness to pay for the different wealth quartiles




Figure 2: Kernel densities associated with estimated willingness to pay for the different types of media exposure




Figure 3: Kernel densities associated with estimated willingness to pay for the different levels of adult female education





Figure 4: Kernel densities associated with estimated willingness to pay for the different levels of adult male education



Figures 5-6 show the wealth and education effects on the WTP²¹. Across each wealth quartile, an additional year of education for female members increases the household's willingness to pay for better quality water more than it does with an additional year of education for male members in the household. Furthermore, moving from no education to 10 years of education (the middle of the range) roughly doubles the expected WTP, which is about the same as the increase resulting from a move from the bottom to the third wealth quartile.

Figure 5: Wealth and female education effects on WTP





Figure 6: Wealth and male education effects on WTP



²¹In these graphs to separate out the effects of the two variables of interest, the values of all other regressors are at their mean levels.

6. Conclusion

Poor environmental quality leads to individuals facing serious health risks in their everyday lives. Individuals will adopt measures to improve their environmental quality only if they perceive the associated health risks and if they can afford to pay for the prevention measure. It is the role of awareness as separate from the income constraint that we try to evaluate in this paper in the context of drinking water in urban India.

Household's awareness about health and hygiene can be raised through formal education and through education imparted (directly and indirectly) by the media. Our results show that there are statistically significant, and quantitatively non-negligible effects of wealth on demand for better quality of environmental goods. However, we also find that there is a strong and comparable effect of education attainment of household members and their exposure to mass media messages through newspapers or radio. This suggests that the common presumption that awareness in comparison to income has a second-order impact on the demand for environmental quality needs to be questioned more generally. Vocal public awareness campaigns to educate the population about the health risks associated with consuming environmental goods of inferior quality may be an important policy instrument.

Acknowledgements

We gratefully acknowledge the financial support provided by the South Asian Network for Development and Environmental Economics (SANDEE). We are grateful to Kirk Hamilton, Rohini Somanathan, and Jeffrey Vincent, participants at the SANDEE workshop and Indian Institute of Management, Kolkata for helpful comments. Dwiraj Bose provided excellent research assistance.

REFERENCES

- Alberini, A., Eskeland, G.S., Krupnick, A., McGranahan, G., 1996. "Determinants of Diarrhoeal Disease in Jakarta." *Water Resources Research*, 32(7): 2259-2269.
- Boyce, J.K., Klemer, A.R., Templet, P.H., Willis, C.E., 1999. "Power distribution, the environment, and public health: A state-level analysis." *Ecological Economics* 29: 127-140.
- Brooks, N. and Sethi, R., 1997. "The Distribution of Pollution: Community Characteristics and Exposure to Air Toxics." *Journal of Environmental Economics and Management*, 32(2): 233-51.
- Colwell, R.R., Huq, A., Islam, M.S., Aziz, K. M. A., Yunus, M., Khan, N.H., Mahmud, A., Sack, R.B., Nair, G.B., Chakraborty, J., Sack, D.A., and Russek-Cohen, E., 2003. "Reduction of cholera in Bangladeshi villages by simple filtration." *Proceedings of the National Academy of Sciences of the USA*, 100: 1051-1055.
- Dasgupta, P., 2001. "Valuing Health Damages from Water Pollution in Urban Delhi, India: A Health Production Function Approach." *Institute of Economic Growth Working Paper Series No. E/210/2001*.
- Filmer, D., Pritchett, L., 2001 "Estimating Wealth Effects without Expenditure Data or Tears: With an Application to Educational Enrollments in States of India." *Demography*, 38(1): 115-132.
- Gadgil, A. 1998. "Drinking Water in Developing Countries." *Annual Review of Energy and Environment*, 23: 253-86.
- Jalan, J., Ravallion, M. 2003 "Does piped water reduce diarrhoea for children in rural India?" *Journal of Econometrics* 112: 153-173.
- Larson, B.A., Gnedenko, E.D. 1999. "Avoiding health risks from drinking water in Moscow: An empirical analysis." *Environment and Development Economics*, 4: 565-581.
- McConnell, K.E., Rosado, M.A., 2000. "Valuing Discrete Improvements in Drinking Water Quality through Revealed Preferences." *Water Resources Research*, 36(6): 1575-82.
- Pargal, S., Wheeler, D., 1996. "Informal Regulation of Industrial Pollution in Developing Countries: Evidence from Indonesia." *The Journal of Political Economy*, 104(6): 1314 - 27.
- Parikh, K.S., J. Parikh, and T.L. Raghu Ram, 1999, Air and water quality management: New initiatives needed in: K.S. Parikh, ed., India Development Report (Oxford University Press, New Delhi).
- World Health Organization, 2002, *The World Health Report 2002*, WHO, Geneva: Switzerland

APPENDIX A

Construction of the wealth index

The wealth index for the i^{th} household is defined as:

$$w_i = \sum_{j=1}^{32} f_j \left[\frac{a_{ij} - m(a_j)}{s_j} \right] \quad \forall i=1, \dots, N$$

In the above equation,

$$\begin{aligned} a_{ij} &= 1 \text{ if } i^{th} \text{ household has asset } a_j, \\ &= 0 \text{ otherwise,} \end{aligned}$$

$$m(a_j) = \frac{\sum_{i=1}^N a_{ij}}{N},$$

$$s_j = \sqrt{\frac{\sum_{i=1}^N a_{ij}^2}{N} - [m(a_j)]^2},$$

and f_j is the “scoring factor” for the j^{th} asset, that is, (f_1, \dots, f_{32}) maximizes the variance of w subject to the constraint $\sum_{j=1}^{32} f_j^2 = 1$.

APPENDIX B

Estimated costs of different purification methods in city of Delhi

Straining: We assume that a household does not have to spend any money if it adopts straining its drinking water with cloth as its chosen water purification method.

Alum tablet: Each alum tablet costs a rupee and can be used to purify 10 liters of water. Assuming a per person daily consumption of drinking water to be 2 liters (this assumption is maintained throughout our cost estimates for the different purification methods), the estimated annual average cost for each person is Rs.73 if the household uses alum tablets as its chosen water purification method.²²

Ordinary filter: Based on a survey of retail shops in Delhi, the average cost price of an ordinary candle filter with a volume capacity of 25 liters is assumed to be Rs.500. In addition, the candles have to be replaced every three years at the cost of Rs.50 for each candle. (Typically these filters come with two candles). Assuming a discount rate of 10 percent and an expected life of the filter to be 10 years, per person annual average cost is estimated to be Rs.110.47. The corresponding estimates under assumptions of 4 and 6 years of expected life of the filter are Rs.276.49 and Rs.237.46 respectively.

Boiling: We assume that a calorie of heat is required to raise the temperature of 1 ml. of water by 1 degree Celsius, the efficiency of a gas stove is 53.6 percent, the calorific value of liquid petroleum gas is 10,600 calories/gram and that there is no heat lost due to improper container. We do not factor in the fixed cost of the gas stove or the container used for boiling in our estimates. Under the above assumptions, if the initial temperature of water that is to be boiled is 25 degrees centigrade, the average annual cost per person of boiling water is estimated to be Rs.1635.20. Under the same assumptions but different initial water temperatures (20 and 30 degrees centigrade), the estimated costs are Rs.1744.70 and Rs.1525.70 respectively.

Electronic filter: Based on a survey of retail shops in Delhi, the average cost price of an electronic candle filter is assumed to be Rs.5,100. In addition, a cost of Rs.250 is incurred towards its maintenance per year for the filter to be effective in killing the possible pathogens in the water. Assuming a discount rate of 10 percent and an expected life of the filter to be 10 years, per person annual average cost is estimated to be Rs.791.53. The corresponding estimates under assumptions of 4 and 6 years of expected life of the filter are Rs.1534.33 and Rs.1116.72 respectively.

²² A daily per capita consumption of 2 liters is the generally accepted value for a person weighing 60 kg. (Gadgil, 1998). This is the value used in estimating ingestion exposure to potentially hazardous chemicals in drinking water. The actual water intake, however, varies considerably from individual to individual, and also according to climate, physical activity and culture.

SANDEE Sponsors



The Beijer International Institute of Ecological Economics –
The Royal Swedish Academy of Sciences

IUCN
The World Conservation Union



FORD FOUNDATION

Education, research, health, sustainability and medicine.
Promote fundamental research and address complex challenges.

MACARTHUR
The John D. and Catherine T. MacArthur Foundation



Swedish International Development
Cooperation Agency

the
abdus salam
international centre for theoretical physics

WORLD BANK INSTITUTE

Promoting knowledge and learning for a better world



**South Asian Network for Development
and Environmental Economics**

SANDEE
P.O. Box 8975 EPC – 1056
Kathmandu, Nepal

Tel: 977-1-552 8761
Fax: 977-1-553 6789
E-mail: info@sandeeonline.org
Website: www.sandeeonline.org